

NOTE

Chromosome constitution of a hybrid between East African Buffalo (*Syncerus caffer caffer*) and Dwarf Forest Buffalo (*Syncerus caffer nanus*)

E. P. CRIBIU and C. P. POPESCU

with the technical collaboration of Jeanine BOSCHER

*Laboratoire de Cytogénétique UNCEIA-I.N.R.A.,
Centre national de Recherches zootechniques, I.N.R.A.,
F 78350 Jouy-en-Josas (France)*

Summary

The chromosome constitution of a male hybrid of first backcross between a female F1 East African × Dwarf African Buffalo (*Syncerus caffer caffer* × *Syncerus caffer nanus*) and a male Dwarf Forest Buffalo (*Syncerus caffer nanus*) had a chromosome number of 53 : 21 acrocentric pairs, 2 unpaired acrocentrics, 3 meta-submetacentric pairs, 1 unpaired submetacentric and 1 acrocentric XY sex chromosome pair. The R-banding pattern shows that the chromosomal difference between the subspecies *caffer* ($2n = 54$) and *nanus* ($2n = 52$) is due to a centric fusion between two acrocentric pairs.

Karyotypes of the two subspecies of African Buffalo, East African (*Syncerus caffer caffer*) and Dwarf Forest (*Syncerus caffer nanus*), have already been described (WURSTER and BENIRSCHKE, 1967; ULBRICH and FISCHER, 1967 and HECK *et al.*, 1968).

The Dwarf Forest type had 54 chromosomes, among which 3 autosomal pairs were meta-submetacentric and 23 were acrocentric, with the X and Y being acrocentric. When compared to the karyotype of Dwarf Forest Buffalo, the East African Buffalo demonstrated 52 chromosomes with one submetacentric pair instead of two acrocentric pairs.

The present report describes the chromosome constitution of a male hybrid of first backcross between a female F1 and a male Dwarf Forest Buffalo. The animal (N° 16569) born in 1969 and kept in Parc Zoologique de Paris, was slaugh-





FIG. 1. — Chromosome constitution of the male hybrid (nanus × (nanus × caffer))

a) Conventional staining karyotype.

b) R-banding karyotype.

Caryotype de l'hybride mâle (nanus × (nanus × caffer))

a) Caryotype en coloration classique.

b) Caryotype avec marquage R.

tered in April 1980. It has recently been included in a study for heredity of coat colour variation by LAUVERGNE and RENVOISÉ (1980).

For chromosome observations, a blood sample was obtained from the jugular vein and cultivated according to the method of DE GROUCHY *et al.* (1964); R-banding was achieved by the technique of DUTRILLAUX *et al.* (1973).

The number of chromosomes in the hybrid is 53, among which a submetacentric and two acrocentrics are not paired. All other chromosome pairs are homomorphic except the XY sex chromosome pair. Two acrocentric pairs exhibit large small arms (fig. 1, *a*).

By using R-banding technique, all chromosome pairs could be identified. R-banding pattern of *p* and *q* arms of the unpaired submetacentric chromosome is characteristic of the two unpaired acrocentric autosomes (fig. 1, *b*).

It is obvious that the unpaired submetacentric chromosome originated from the East African type whereas the two unpaired acrocentric autosomes were derived from the Dwarf Forest type.

In regard to the karyotype evolution of the two subspecies of African Buffalo, it could be suggested that the Dwarf type with $2n = 54$ occurred first and during evolution, the East African type should have developed by Robertsonian fusion of two acrocentric pairs.

Received for publication in november 1980.

Résumé

Le caryotype d'un hybride entre le Buffle Africain (Syncerus caffer caffer) et le Buffle nain (Syncerus caffer nanus)

Le caryotype d'un mâle, produit de croisement de retour d'une femelle hybride F₁ Buffle Africain × Buffle nain (*Syncerus caffer caffer* × *Syncerus caffer nanus*) et d'un Buffle nain mâle (*Syncerus caffer nanus*) est composé de 53 chromosomes : 21 paires acrocentriques, 2 chromosomes acrocentriques non appariés, 3 paires méta-submétacentriques, 1 chromosome submétacentrique et 2 chromosomes sexuels X et Y acrocentriques. Le marquage R a montré que la différence de caryotype de ces deux sous-espèce est due à une fusion centrique entre deux paires de chromosomes acrocentriques.

References

- DE GROUCHY J., ROUBIN M., PASSAGE E., 1964. Microtechnique pour l'étude des chromosomes humains à partir d'une culture de leucocytes sanguins. *Ann. Génét.*, **7**, 45.
- DUTRILLAUX B., LAURENT M., COUTURIER J., LEJEUNE J., 1973. Coloration pour l'acridine orange de chromosome préalablement traités par le 5 bromodeoxyuridine (B.U.D.R.). *C. R. Acad. Sci. (Paris)*, **276**, 3179-3181.
- HECK H. D., WURSTER D. H., BENIRSCHKE K., 1968. Chromosome study of numbers of the subfamilies Caprinae and Bovidae: the muskox, ibex, aoudad, Congo buffalo and Gaur. *Z. Säugetierk.*, **33**, 172-179.
- LAUVERGNE J. J., RENVOISÉ C., 1980. Héritéité des variations de couleur du pelage du Buffle africain (*Syncerus caffer*). *Ann. Génét. Sél. Anim.*, **12**, 1-7.
- ULBRICH F., FISCHER M., 1967. The chromosomes of the Asiatic buffalo (*Bubalus bubalis*) and the African buffalo (*Syncerus caffer*). *Z. Tierz. Züchtbiol.*, **83**, 219-223.
- WURSTER D. H., BENIRSCHKE K., 1967. The chromosomes of twenty-three species of *Cervoidae* and *Bovidae*. *Mammalian Chromosomes Newsletter*, **8**, 226-229.