

registered and treated as all-or-none traits in genetics analyses. A significant sire effect was found for all serum traits except albumin, with heritability estimates between 0.1 and 0.2. The disease characters showed heritability estimates between 0.04 and 0.08. Immunoglobulin and total-protein concentrations seemed to have a negative genetic correlation to diseases.

Genetic polymorphisms of plasma α_1 -protease inhibitors in some domestic animals

R. KUMAR JUNETA and B. GAHNE

*Department of Animal Breeding and Genetics,
Swedish University of Agricultural Sciences, S-750 07 Uppsala, Sweden*

The genetic polymorphism and phenotypes of plasma α_1 -protease inhibitor (α_1 -PI) in horse, cattle, sheep, goat, pig, dog and mink were described in a brief review. In all these species, two α_1 -protease inhibitors (Pi-1 and Pi-2) were observed. Pi-1 showed inhibition for both trypsin and chymotrypsin while Pi-2 showed inhibition for only trypsin. In horse 6 Pi-1 and 5 Pi-2 alleles were observed. There was a very close genetic linkage between Pi-1 and Pi-2 loci in horse. The horse Pi-1 and Pi-2 fractions together correspond to the horse acidic prealbumin (Pr) fractions described earlier in literature.

In cattle, sheep and goat, polymorphism was observed for only Pi-2. Three alleles in cattle, three in sheep and two alleles in goat were observed.

In pig, the earlier described polymorphism pre-albumin (Pa) was identified as α_1 -protease inhibitor. Two Pa alleles have been reported in pig.

In dog and mink, polymorphism was observed only for Pi-1. Three Pi-1 alleles were observed in several breeds of dogs. In mink, the polymorphic post-albumin (Pa) described in literature, was identified as α_1 -protease inhibitor.

The α_1 -PI fractions in all these species are present in high concentration in plasma and thus could be visualised by general protein staining of gels after electrophoresis. The possible association of α_1 -PI phenotypes with respiratory and inflammatory diseases in domestic animals was also briefly discussed.

Immune response to some *E. coli* antigens in swine

I. EDFORS-LILJA *, B. GAHNE *, B. MOREIN ** and H. PETERSSON *

** Dep. of Animal Breeding and Genetics, Swedish University of Agricultural Science,
S-750 07 Uppsala, Sweden*

*** National Veterinary Institute, S-104 05 Stockholm, Present address :
Dep. of Virology, Swedish University of Agricultural Science, Box 585
S-751 23 Uppsala, Sweden*

The humoral immune responses to two *E. coli* antigens, K88 and 0149, were studied in 60 pigs after 9 sires and from 19 litters. One animal per litter was in addition kept as a control. The animals were immunized subcutaneously with a whole cell suspension of *E. coli* at the age of approximately 10 weeks. Ten of the animals were given a second immunization 3 weeks after the first one. Blood samples were taken immediately before immunization, 1 and 2 weeks post-injection. The total amount of specific antibodies in serum to K88 and 0149 antigen were analysed by ELISA technique. A significant increase in antibody titers was obtained after immunization, although a pronounced individual variation was seen. The animals were given a constant dose of the suspension without regard to