

## R-banding pattern of the prometaphase chromosomes of the domestic sheep *Ovis aries* L.

D. Di Berardino <sup>1</sup>, M.B. Lioi <sup>2</sup>, C. Miranda <sup>1</sup>, A. Di Milia <sup>1</sup>,  
M.G. D'Agostino <sup>1</sup> and D. Matassino <sup>1</sup>

<sup>1</sup> University of Naples, Department of Animal Production, 80055 Portici, Naples

<sup>2</sup> University of Basilicata, Institute of Animal Production, 85100, Potenza, Italy

(received 19-2-1988, accepted 5-5-1988)

**Summary** — The RBA-banding pattern of the domestic sheep *Ovis aries* L. and the diagrammatic representation at the 510 band stage are presented and proposed as the "standard" RBA-karyotype for this species.

**domestic sheep – RBA-banding – chromosomes**

**Résumé** — Description des chromosomes marqués en bandes R du mouton domestique *Ovis aries* L. Les bandes R (technique RBA) des chromosomes du mouton domestique et son idiogramme à l'état de 510 bandes sont présentés et proposés comme le caryotype «standard» de cette espèce.

**mouton domestique – bandes R – chromosomes**

### Introduction

The discovery of the RBA-banding procedure by Dutrillaux *et al.* (1973) has provided a remarkable improvement in the identification and description of individual chromosomes of mammals, including man. The benefits of this technique are quite evident in the family Bovidae, whose chromosomes, especially the smallest elements of the karyotype, are not easily distinguishable when G-banded; the difficulty lies in the fact that in Bovidae chromosomes, both centromeres and telomeres are mostly G-negative and, therefore, the smallest autosomes have to be identified by relying upon very few and often undefined bands.

The potential of the RBA-banding technique for definite identification of Bovidae chromosomes, first pointed out by Popescu (1975) and by Gustavsson and Hagelthorn (1976), has subsequently been demonstrated by another series of studies on the R-banding pattern of prometaphase chromosomes of *Bos taurus* L. (Di Berardino *et al.*, 1979, 1985; Di Berardino and Iannuzzi, 1982), *Bubalus bubalis* L. (Di Berardino *et al.*, 1981; Di Berardino and Iannuzzi, 1981, 1984) and *Capra hircus* L. (Di Berardino *et al.*, 1987).

As a contribution to establishment of the standard RBA-banded karyotype of *Ovis aries* L., this paper presents karyotypes and diagrammatic representations of the RBA-banding patterns at the 510 band stage.

## Materials and Methods

Peripheral blood drawn from the jugular vein of 11 animals of the Gentile di Puglia breed (6 males and 5 females) was cultured for 72 h in RPMI 1640 medium (Flow, Dutch modification), supplemented with 10% fetal calf serum (Gibco), antibacterial and antifungal agents, 0.1% of L-glutamine, and pokeweed mitogen (Gibco). After 48 h, lymphocytes were synchronized by adding excess thymidine (0.3 mg/ml, final concentration). The S-phase block was released 18 h later by washing the cultures in fresh RPMI medium. To induce R-banding the cells were recultured in growth medium as described above, supplemented with BrdU (Sigma, 10 µg/ml, final conc.) added 1 h later. The optimal recovery time was found to be around 6.5 h, including 1 h of exposure to colcemid solution (Gibco).

After hypotonic treatment with 0.075 M KCl for 20 min at 37.5°C, the cells were fixed in methanol-acetic acid 3:1 for 1 h, centrifuged, fixed again, and left overnight in a refrigerator. The next day the cell suspension was centrifuged, fixed again in fresh fixative, spread on to clean wet slides and air dried.

The staining procedure for RBA-banding was performed as described by Di Berardino and Iannuzzi (1982).

## Results

Fig. 1 shows a representative prometaphase RBA-banded karyotype of the sheep ( $2n = 54,XY$ ).

The karyotype has been arranged by following the same nomenclature as previously adopted for the RBA-banded karyotype of the goat (Di Berardino *et al.*, 1987) and cattle (Di Berardino *et al.*, 1985) which refers, as far as possible, to the Reading system (Ford *et al.*, 1980).

Several karyotypes were prepared from the 11 investigated animals, but only the best banded chromosomes, 4 for each pair, were selected and used to produce the diagrammatic representation shown in Fig. 2A, B, C.


The whole idiogram of the RBA-banding pattern of sheep chromosomes is presented in Fig. 3A, B. The numbering of the bands within each individual chromosomes follows the ISCN nomenclature (ISCN, 1978).

The RBA-banding pattern of the individual sheep chromosomes has been found to be identical to that of the goat chromosomes previously reported (Di Berardino *et al.*, 1987); therefore, the detailed description is omitted.


Given the complete homology in banding pattern between the chromosomes of the 2 species, and following a phylogenic criterion, the goat chromosome nomenclature, which is identical to that of cattle, was adopted to classify the sheep chromosomes.

For this reason there is no correspondence between the G-banded standard karyotype presented by Long (1985) and the present one.

Table I shows the nomenclature of the corresponding homologous chromosomes between sheep and goat; the 3 pairs of submetacentric chromosomes in the sheep karyoty-



**Fig. 1.** Representative RBA-banded karyotype of the domestic sheep ( $2n = 54,XY$ ), proposed as standard.



**Fig. 2A,B,C.** RBA-banding pattern of individual prometaphase chromosomes of the domestic sheep. Groups of 4 representative chromosomes, selected from different karyotypes, were used to construct the diagrammatic RBA-banding pattern shown on the left of each group.




Fig. 2B. Legend as in Fig. 2A.





Fig. 2C. Legend as in Fig. 2A.




1

2

3




4




5



6



7



8

Fig. 3A,B. Diagrammatic representation of the RBA-banding pattern of prometaphase chromosomes of sheep.




Fig. 3B. Legend as in Fig. 3A.

**Table I.** Corresponding homologous chromosomes in the sheep and goat RBA-banded karyotypes.

Sheep . . . .	1p	q	2p	q	3p	q	4	5	6	7	8	9	10	11	12	13
Goat . . . . .	3	1	8	2	11	5	4	6	7	9	10	12	13	14	15	16
Sheep . . . . .	14	15	16	17	18	19	20	21	22	23	24	25	26	X	Y	
Goat . . . . .	17	18	19	20	21	22	23	24	25	26	27	28	29	X	Y	

pe are the results of 3 different sets of centric fusions involving, respectively, chromosomes 1-3, 2-8, and 5-11 of the goat karyotype.

The total number of bands counted in the haploid set of sheep chromosomes in the prometaphase stage, including the X and Y chromosomes, equalled 510 bands, of which 187 (36.7%) were positive, 251 (49.4%) negative, and 72 (14.1%) intermediate.

## Discussion and Conclusion

The present paper has to be considered as a contribution to the establishment of the standard RBA-banded karyotype of the species *Ovis aries* L.

Lymphocytes were synchronized by excess of thymidine and RBA-banding was induced by using a combination of BrdU and H33258, as previously reported for the R-banded karyotype of the goat (Di Berardino *et al.*, 1987). Thymidine synchronization is achieved by a feed-back inhibition mechanism which blocks the formation of deoxycytidine triphosphate from cytidine 5-phosphate (Xeros, 1962), thus blocking the majority of the cells at mid S-phase of the cell cycle. Thymidine was preferred to methotrexate or fluorouracil as it is less toxic and less clastogenic for cells (Dutrillaux and Viegas-Pequignot, 1981; Ronne *et al.*, 1984; Ronne, 1984).

A remarkable increase in the R-band resolution was achieved by combining with BrdU H33258, which links to DNA by hydrophobic bonds (Bontemps *et al.*, 1975; Comings, 1975) with a 4-fold stronger affinity to poly (dA-dbrdU) than poly (dA-dT) segments (Latt and Wohlleb, 1975); this combined incorporation delays chromosome contraction and enhances band contrast (Ronne, 1983).

The present RBA-banded karyotype can be utilized for further studies on detection of numerical as well as structural chromosomal abnormalities, evolutionary relationships among the members of the family Bovidae, and gene mapping by using *in situ* hybridization procedures (Geffroin *et al.*, 1984). The extensive degree of banding homology among the chromosomes of sheep, goat, buffalo, and cattle may be helpful for the localization of common genes responsible for important economic traits in animal production.

## Acknowledgments

This work has been supported by a financial contribution from the National Research Council, Rome, Italy.

## References

- Bontemps J., Houssier C. & Fredericz E. (1975) Physicochemical study of the complexes of 33258 Hoechst with DNA and nucleohistone. *Nucleic Acids Res.* 2, 971-984
- Comings D.E. (1975) Mechanisms of chromosome banding. VIII. Hoechst 33258-DNA interactions. *Chromosoma*, 52, 229-243
- Di Berardino D. & Iannuzzi L. (1981) Chromosome banding homologies in Swamp and Murrah buffalo. *J. Hered.* 72, 183-188
- Di Berardino D. & Iannuzzi L. (1982) Detailed description of R-banded bovine chromosomes. *J. Hered.*, 73, 434-438
- Di Berardino D. & Iannuzzi L. (1984) Detailed description of RBA-banded chromosomes of river buffalo (*Bubalus bubalis* L.). *Génét. Sé. Evol.*, 16, 249-260
- Di Berardino D., Iannuzzi L. & Lioi M.B. (1985) The high resolution RBA-banding pattern of bovine chromosomes. *Cytogenet. Cell Genet.* 39, 136-139
- Di Berardino D., Iannuzzi L., Ferrara L. & Matassino D. (1979) A new case of Robertsonian translocation in cattle. *J. Hered.* 70, 436-438
- Di Berardino D., Iannuzzi L., Bettini T.M. & Matasino D. (1981) Ag-NORs variation and banding homologies in two species of Bovidae: *Bubalus bubalis* and *Bos taurus*. *Can. J. Genet. Cytol.* 23, 89-99
- Di Berardino D., Ronne M., Burgete I., Lioi M.B., Taibi L. & Matassino D. (1987) The R-banding pattern of the prometaphase chromosomes of the goat (*Capra hircus* L.). *J. Hered.* 78, 225-230
- Dutrillaux B. & Viegas-Pequignot E. (1981) High resolution R- and G-banding on the same preparation. *Hum. Genet.* 57, 93-95
- Dutrillaux B., Laurent C., Couturier J. & Lejeune J. (1973) Coloration des chromosomes humains par l'acidine orange après traitement par la 5-bromodeoxyuridine. *C. R. Acad. Sci. Sér. D.* 276, 3179-3182
- Ford C.E., Pollock D.L. & Gustavsson I. (1980) Proceedings of the 1st International Conference for the Standardisation of Banded Karyotypes of Domestic Animals. *Hereditas* 92, 145-162
- Geffrotin C., Popescu C.P., Cribeu E.P., Boscher J., Renard C., Chardon P. & Vaiman M. (1984) Assignment of MHC in swine to chromosome 7 by *in situ* hybridization and serological typing. *Ann. Génét.* 27, 213-219
- Gustavsson I. & Hagelthorn M. (1976) Staining technique for definite identification of individual cattle chromosomes in routine analysis. *J. Hered.* 67, 175-178
- ISCN (1978) An international system for human cytogenetic nomenclature. *Cytogenet. Cell Genet.* 21, 309-404
- Latt S.A. & Wohlleb J.C. (1975) Optical studies of the interaction of 33258 Hoechst with DNA, chromatin and metaphase chromosomes. *Chromosoma* 52, 297-316
- Long S.E. (1985) Standard nomenclature for the G-banded karyotype of the domestic sheep (*Ovis aries* L.). *Hereditas* 103, 165-170
- Popescu C.P. (1975) Essai d'identification des chromosomes bovins (*Bos taurus*) à l'aide du marquage au 5-bromodeoxyuridine (BrdU). In : 2<sup>e</sup> Colloque Européen de Cytogénétique des Animaux Domestiques, Giessen, 29-30 septembre 1975. INRA-CNRS, France, pp. 59-64
- Ronne M. (1983) Simultaneous R-banding and localization of dA-dT clusters in human chromosomes. *Hereditas* 98, 271-278
- Ronne M. (1984) Fluorouracil synchronization of human lymphocyte cultures. Induction of high resolution R-banding by simultaneous *in vitro* exposure to 5-bromodeoxyuridine/ Hoechst 33258. *Hereditas* 101, 215-208
- Xeros N. (1962) Deoxyriboside control and synchronization of mitosis. *Nature* 194, 682-683