

# Association of the porcine *C3* gene with haemolytic complement activity in the pig

Supamit MEKCHAY, Siriluck PONSUKSILI,  
Karl SCHELLANDER, Klaus WIMMERS\*

Institute of Animal Breeding Science, University of Bonn,  
Endenicher Allee 15, 53115 Bonn, Germany

(Accepted 4 February 2003)

**Abstract** – The complement component *C3* plays an essential role in the activated complement system, which is involved in phagocytosis, inflammation and immunoregulation to destroy infectious microorganisms. The *C3* molecule has more implications in the general defence mechanisms. In this study, the porcine *C3* cDNA sequences including 5'- and 3'- flanking regions were determined and the polymorphisms in this gene were identified to carry out an association analysis between *C3* and complement activity traits. Porcine *C3* gene has high homology with human *C3*. Five single nucleotide polymorphisms (SNPs) and one microsatellite were detected in the porcine *C3* gene. Haemolytic complement activity of alternative and classical pathways (ACH, CCP) was measured in 416 F2 animals of a crossbred of Duroc × Berlin Miniature Pig, which were immunized with *Mycoplasma*, Aujeszky and PRRS vaccines. *C3* markers were found to be significantly associated ( $P < 0.05$ ) with both ACP and CCP. Animals with the more frequent haplotype present in Duroc and other commercial breeds exhibit higher ACP and CCP levels than the animals with haplotype specific to some Berlin Miniature Pigs. The association of *C3* with complement activity reinforces the importance of *C3* as a candidate gene for natural resistance to microorganisms.

**porcine *C3* gene / natural resistance / haemolytic complement activity / association / pig**

## 1. INTRODUCTION

The complement cascade defines an important link between the innate and the specific immune system [13]. The complement component *C3* is the key molecule of the three pathways of complement activation (alternative, classical and lectin pathways), which are involved in phagocytosis, inflammation and immunoregulation processes to destroy infectious microorganism [19].

---

\* Correspondence and reprints  
E-mail: kwim@itz.uni-bonn.de

*C3* deficiency in hosts causes increasing susceptibility to bacterial or viral infections [1,4,13,22]. *C3* has more implication in general defence mechanisms. We have previously found polymorphic sites within the porcine *C3* which were used for genetic mapping of the locus [25]. The present study was carried out to investigate the porcine *C3* gene as a candidate gene for complement activity traits. Therefore, the full length cDNA sequence of the porcine *C3* as well as 5'- and 3'- flanking regions were obtained, screened for additional polymorphisms in this gene and, finally, association analysis was conducted between *C3* markers and complement activity traits.

## 2. MATERIALS AND METHODS

### 2.1. Animals


Sequence information of the porcine *C3* gene was identified in a DUMI F2-animal of a reciprocal cross of Duroc × Berlin Miniature Pig [9]. Screening for polymorphisms in the porcine *C3* gene was performed in Duroc (n = 6), German Landrace (n = 28), Pietrain (n = 23), DUMI F2 (n = 902) and Thai native (n = 32) pigs. For association study, haemolytic complement activity was recorded in 416 animals of the DUMI-F2 resource population.

### 2.2. cDNA sequence and 5' and 3' flanking regions of the porcine *C3* gene

Total RNA was extracted from pig liver using TRIzol<sup>TM</sup> Reagent (Life Technologies, Karlsruhe, Germany). Full length porcine *C3* cDNA sequence was determined by the SMART<sup>TM</sup> RACE cDNA Amplification Kit (Clontech, Heidelberg, Germany). RACE primers and gene specific primers were designed based on partial porcine *C3* sequence information (GenBank accession number AF110278, F14640) (Fig. 1):

- RACE: 5'-ctaatacgactcactatagggcaagcagtggtatcaacgcagagt-3'
- C3-A2: 5'-ccttctccacgacatcccagatcctacg-3'
- C3-B1: 5'-ccaccaagaccatgaatgtg-3'
- C3-B2: 5'-tagagcttctggccaggttc-3'
- C3-C1: 5'-ggtggtgacatgtaccacg-3'.

The 5' and 3' flanking regions of porcine *C3* gene were isolated from a genomic DNA library, which was constructed using Lambda DASH II<sup>®</sup> (Stratagene, Amsterdam, Netherlands). Two positive clones of 5' and 3' flanking region were amplified with primers specific to porcine *C3* (5'-flanking region 5'-tgtgcccttctctgatttg-3', 3'-flanking region 5'-gctccgatgaagtgcaggt-3') and T3/T7



**Figure 1.** Amplification of the full-length porcine C3 gene. Three PCR fragments (A, B and C) were amplified with primers based on available porcine C3 sequence in GenBank, a 580-bp partial cDNA fragment (shaded area 1) (accession No. AF110278) and a 462-bp EST of porcine C3 (shaded area 2) (accession No. F14640).

standard primers. All the PCR fragments were ligated in pGEM<sup>®</sup>-T vector (Promega, Mannheim, Germany) and sequenced using the SequiTherm Excel cycle sequencing Kit (Epicentre Technologies, Biozym, Hessisch Oldendorf, Germany) and a LI-COR 4200 automated sequencer (LI-COR Biosciences, Bad Homburg, Germany).

### 2.3. Polymorphism screening and marker genotyping

To identify polymorphisms in the porcine C3 gene, oligonucleotide primers were designed revealing 17 overlapping PCR fragments (Tab. I). They were amplified from individual liver cDNA and genomic DNA of six pig breeds. PCR was performed in a 25  $\mu$ L reaction volume containing 50 ng of liver cDNA, 0.2  $\mu$ M of each primers, 50  $\mu$ M each dNTPs, 0.5 U of *Pfu* polymerase (Promega) in 1 $\times$  *Pfu*-PCR buffer. PCR cycling program was 94  $^{\circ}$ C for 3 min, followed by 35 cycles of 94  $^{\circ}$ C for 30 s, 52–60  $^{\circ}$ C (Tab. I) for 30 s, 72  $^{\circ}$ C for 1 min and final extension at 72  $^{\circ}$ C for 5 min. PCR products were gel purified, cloned and sequenced. The nucleotide sequences of each animal were compared to find out polymorphisms in the porcine C3 gene.

To genotype each polymorphic site in the porcine C3 gene, PCR-RFLP, allele specific PCR, SSCP and microsatellite analysis were performed. Oligonucleotide primers (Tab. II) were used to amplify PCR fragments from genomic DNA. PCR reactions were carried out in 20  $\mu$ L volume using *Taq* polymerase (Promega). Amplified PCR fragments with primer C3-I (Tab. II) were mixed with 1:10 with loading buffer (95% formamide, 10 mM NaOH, 0.25% bromophenol blue, 0.25% xylene cyanol), denatured at 95  $^{\circ}$ C for 5 min, then cooled on ice immediately and separated on 12% polyacrylamide gel (49:1 acrylamide/bis-arylamide) at room temperature with a constant 12 W for 6 h in 0.5 $\times$  TBE. The SSCP-bands were stained by silver staining procedure. Amplicon C3-II was digested with restriction enzyme *TaqI* (Promega). The digested PCR fragments were analysed on 2% agarose gels. Genotyping the SNPs within amplicons C3-III and C3-IV was done as described previously [25]. Fragments with length polymorphism were amplified with primer C3-V (Tab. II) and were identified on 6% SequiGel<sup>®</sup>XR-denaturing polyacrylamide gels

**Table I.** Primer sequences used for screening polymorphisms in the porcine *C3* gene.

Name	Forward primer sequence (5' to 3')	Reverse primer sequence (5' to 3')	Annealing temp. (°C)	cDNA (bp)	Genomic- DNA (bp)
C3-5'	GCATCGACTTGAATTCACAG	CCATGATGGAGGGACGGTG	55	–	477
C3-01	TCCTTTCCCTCTGTCCCTTT	GGGAGTCCCGTTTGATGTC	58	577	–
C3-02	TGAGACACCTGAAGGCATTG	GCTGTTCCATCCACACTCTC	58	306	~ 700
C3-03	TCAACATCATTGCCAGGTTC	GGGTCTTGGTGAAGTGGATC	59	324	–
C3-04	CATCGTGACCTCCCCCTATC	GTGCGCAGGTGGAAGTTAAC	59	393	~ 1000
C3-05	TCAAGCCAGGGGAGAATCTC	TCTCCAGGGTCATCTGTTGC	56	360	~ 1500
C3-06	GAAGCAAGACAAGCAGCATC	TGTCCATCCTTTTCTCCATG	56	338	~ 900
C3-07	GTGCAGCTCATGGAGAAAAG	ATTCATGGTCTTGGTGGAG	56	360	–
C3-08	CTCCACCAAGACCATGAAT	TCTTCAAGGGCACAATG	54	356	–
C3-09	CATTGTGCCCTTGAAGA	GTGCTGTCCAGGTAGTGC	55	396	–
C3-10	CAGTCATCGCTGTGCACTA	GTTCTTGAAGCCACCAATC	55	343	–
C3-11	GATTGGTGGCTTCAAGAAC	CTACCAGCAGAGCCAAG	52	341	–
C3-12	CTTGGCTCTGCTGGTAG	CGTGGTACATGGTCAC	52	338	~ 1000
C3-13	GTGACCATGTACCACG	CCTCCAGGGTGTGTGAGAC	52	337	~ 1300
C3-14	GATCTCACACACCCTGGAGG	CACTTCATCGGAGCCTGATT	55	379	–
C3-15	AATCAGGCTCCGATGAAGTG	GGCAACCAAAGACGACCAT	56	270	–
C3-3'	TGGCCAACTTCTCTGAGAAC	TGCAGGCAGATACAGTTGAG	60	–	530

**Table II.** Nucleotide sequence primers for genotyping the polymorphisms in the porcine *C3* gene.

Amplicon	Primer (5' to 3')	Size (bp)	Annealing temp. (°C)
C3-I	F: GCATCGACTTGAATTCACAG	477	55
	R: CCATGATGGAGGGACGGTG		
C3-II	F: CACCCTGATTGCTGCCAATG	383	60
	R: TACCTCAACTTACTGCGGTC		
C3-III	F: TGAGAATGTGGATGGACCAG	384	60
	R: GGACTTGAATGCCCAAGATC		
C3-IV	F1: AAGGATCTGAACCTGGATGTA	454	64
	F2: GGATCTGAACCTGGATGTG	452	
	R: ACCCCGCTAATCTGTGATGC		
C3-V	F: TGGCCAACTTCTCTGAGAAC	204, 207	60
	R: TAGGATGAACCTGAGCTGTG	211	


(Biozym Diagnostik GmbH) by the LI-COR 4200 automated sequencer (LI-COR Biosciences). The fragment size of alleles was analysed by One-Dscan software (Scanalytics, MWG Biotech).

#### 2.4. Haemolytic complement activity phenotypes

Total haemolytic complement activities were determined in 416 F2-animals of the DUMI-resource population produced from 11 F1 sows (2 full-sib groups) mated with 3 F1 boars. The animals were reared in the Frankenforst research farm of the University of Bonn. They were immunized with Mycoplasma, Aujeszky and PRSS vaccines at 6, 14 and 16 weeks of age, respectively. Blood samples were taken immediately before vaccination and on 4 and/or 10 days after vaccination (Fig. 2). Total haemolytic activity of the alternative (ACP) and classical complement pathway (CCP) was determined by a method modified from Liu and Young [15]. Non sensitised rabbit erythrocytes and sensitised sheep erythrocytes were used as target cells, for ACP and CCP respectively. The haemolytic complement activity was expressed as the titre that lysed 50% of erythrocytes (CH50 unit · mL<sup>-1</sup>).

#### 2.5. Association analysis between *C3* and complement activity traits

Association between *C3* markers and haemolytic complement activity traits was analysed using the REPEATED statement of the SAS<sup>®</sup> PROC MIXED procedure [14]. The statistical model included *C3* genotype, time of blood



**Figure 2.** Vaccination programme and time of blood sampling in the DUMI-F2 resource population.

sampling, sire, and interaction of *C3* genotype and time of blood sampling as fixed effects, and dam nested within sire as random effect. A heterogeneous compound-symmetry (CSH) covariance structure was included in the statistical model to analyse fixed effects because it gave the best fit when compared to other covariance structures.

### 3. RESULTS

#### 3.1. Sequence analysis of the porcine *C3* gene

The complete porcine *C3* cDNA sequence (accession No. AF154933) was determined. The 5127 bp cDNA contains an open reading frame coding for 1661 amino acids, including a 22-amino acids signal peptide, 643 amino acids  $\beta$ -chain, a 4 amino acids linker and 992 amino acids  $\alpha$ -chain. The ORF of porcine *C3* shows 80, 78 and 77% homology with human, rat and mouse *C3* nucleotide sequence, respectively [6,7,17]. The 63 bp 5'-UTR is a pyrimidine rich region (approximately 80% pyrimidines). A 56 bp of 3'-UTR was sequenced and a polyadenylation signal (AATAAA) was found at position 5074 to 5079 of cDNA sequence (AF154933). The sequence information of the 5'-flanking region (Fig. 3a) contained several features of promoter region. The putative TATA-box (GATAAA) sequence was located at -29 bp of the first nucleotide *C3* sequence (AF154933) and various putative enhancer/transcription factor binding sequences were also found (Tab. III). Comparison of the porcine 5'-flanking sequence with the promoter region of human *C3* sequence (X62904) [10] revealed 74% sequence identity. However, the 3'-flanking region (Fig. 3b) of porcine *C3* had low homology with human *C3* sequence (M63422) [21].

#### 3.2. Polymorphism and allelic frequency of the porcine *C3* gene

Six polymorphic sites were found in the porcine *C3* gene including five SNPs and one microsatellite (Fig. 4a). Two SNPs were found in the 5'-flanking

```

-487  GACTCTTGCT  GCCAGCATCG  ACTTGAATTC  ACAGGAAGGG  TTTGGTTAAT
-437  AAGTGCATGG  CCAAAAGTGG  ACTGAGAGCC  AGGTGCAGAT  ATCGGGGAAG
-387  GGGGGCAGGC  ACTATCTGGG  GAAGAGAGAA  TAATTTTCCT  CCACTACAAA
-337  GTGGGTATAG  CAGGGACAGG  TCCCCATTCA  CCAGCCAGCC  TTGACCATGA
-287  GAGAGGCCAG  GGCAGGAGGC  TGTTGTATT  TTGTTGCAGG  AGGACGTGCC
-237  CTTAGGTTAT  TTTCCCAGG  ACTGTGATGG  CCACACAGAT  TGAAAACTT
-187  TGGAAATGAC  ATTGAGAAAT  CTGGGGCAGC  CCCAGGGAGG  GGGGAGGCCA
-137  CAGGGAGTGG  AGGGCTGGGC  TGAAGAGGGG  GAAAAGCAGC  TGCCAGATAA
-087  AAGGCAGCCT  CCAGCAGCCT  CTGCTCACTT  CCCCCCCAC  CCCCGTCCCTT
-037  TCCCTCTGTC  CCTTTGTCC  TCCACCGTCC  TCCCATCATG  .....
...  C3 cDNA sequence in GenBank accession F154933  ...
1    TGATGCCACT  CCCCCACAGT  CTACCCAATA  AAGCTCCAGT  TATCTTTCCAC
51   ATTTCCCCCC  ACATCTGAAG  GTCTTTTTTT  TTTTTTTTTT  GAAGGTCTTT
101  GAGAGGGGAA  AAGAGGCAGC  TGTGATCCCG  CCACTGCCTC  ACCCACAGCT
151  CAGGTTTCATC  CTACTGAAAC  TCCACCTGCT  TCCCACTTCA  TCCCCTCCAG
201  CAGTTCACCC  GGCTTCCTGG  CCTCAAGTGC  ATGTACCTGC  CTTACCTGCA
251  CCTAGCGGGC  AGTACATCTG  CAGCTTCCCC  TGCATCTCCC  CTGCACCCC
301  ATCCTTACCC  AAATTTCAT  TTACATTCTC  ACCTGCATCC  TCCCCAGCAG
351  CTCACTGTCC  CTTACCTGA  GCTTTCACCA  CCTTGCTTAC  CTACCTCCTT
401  ATTTTCATCCT  CACCTGAGCT  TTTCCCAATA  GTCTCAGCTG  AATCTTCACC
451  CTCTCTCACC  TGCACACTCA  ACTGTATCTG  CCTGCATTTT  GTGATCTTCC
501  CCTGCTTCCC  CCACCTACAT  CTTAATCTGT  AACTTAACT  CTGACTCTAC
551  CAACATCTTT  TCCTTTATTC  TCACCTTGTC  TTTCTACCTG  TATCCTTACC
601  TGTATTTTCT  CTGCATCCTC  CTCAGGGTGA  GGAAGTGCAT  CTTTGCCTAC
651  ATCCCTCTCT  AGTCTCTTTC  CTGTGTTCTC  ACCAGCATCC  TTAGCTACAT
701  TCTCACCTAC  ATCTTCCCCT  GCATCTTTTT  TTTGCTTTTT  TTTTTTTTGC
751  TTTTTTTTTG  CTTTTTAGGG  CCACACCCAC  GGCATATGGA  GGTTCCCAAG
801  CTAGGGGGCA  AATTGGAGCT  GCAGCTGCCA  GCCTCACCAC  AGCCACATCA
851  ATGCAGGATC  CGAATCTTTC  GCCCTATAGT  GAGT

```

**Figure 3.** Sequence of the 5'-flanking region upstream from ATG start codon (a) and of the 3'-flanking region downstream from TGA codon (b) of the porcine C3 gene (accession No. F154933). The 5'-UTR is indicated in italics and the putative transcription start site is indicated in italics, bold and underlined. A putative TATA box and polyadenylation signal are indicated in bold and underlined in the 5'- and 3'-flanking regions, respectively.

region at position -384 (G > A) and -44 (C > T) nucleotide (nt) upstream ATG start codon. The other three SNPs were located in the coding region at 1905 (C > A), 3882 (G > A) and the nt 204 (T > C) of the intron 13. A microsatellite with (T)<sub>14-21</sub>-repeat units was found in the 3'-flanking region of C3 and was located at 71 bp downstream from the TGA stop codon.

The two SNPs of 5'-flanking region were genotyped by SSCP analysis (Fig. 4b). *Taq* I-RFLP was used to detect the variation of the nt 204 in the intron 13 of C3-cDNA sequence (amplicon C3-II) and showed either undigested PCR product (384 bp, allele T), and/or digested product (285 bp and 143 bp, allele C) (Fig. 4c). Similarly, restriction enzyme *Hsp92* I was used to test polymorphism at position 1502 (amplicon C3-III) and revealed either

**Table III.** Position of transcription factor binding sites in the 5'-flanking region of the porcine *C3* gene.

Factor	Consensus sequence	Position	Homology (%)	Ref.
Enhancer core	GGGGAAAA	-110 to -103	88	[8]
C/EBP	TTGAGAAAT	-176 to -168	100	[23]
IL-6-RE	AGGGGGA	-112 to -106	86	[21]
	TCTGGGG	-168 to -162	100	[21]
	TGAGAAA	-175 to -169	100	[21]
	TGAAAAA	-197 to -191	100	[21]
	TGAGAGA	-290 to -284	100	[21]
	TGGGGAA	-371 to -365	100	[21]
IL-6 RE rev	TTCCCCA	-220 to -226	100	[8]
	TCCTCCA	-351 to -345	100	[8]
IFN $\gamma$ RE	GAAAACTTTGGAAA	-196 to -192	87	[8]
AP-2	CCCTTAGG	-132 to -139	88	[18]
LF-A1	TGGCCA	-210 to -205	83	[8]
	TGGACT	-420 to -415	83	[8]
LF-B1/HNF-1	GTTAAT	-443 to -438	100	[8]
	GTTATT	-232 to -227	83	[8]
Estrogen RE	TGTCCTTTGTCC	-31 to -19	85	[21]


digested PCR product (237 bp and 146 bp, allele C) and/or 383 bp (allele A) of undigested product (Fig. 4d). The SNP located at 3882 was genotyped by allele specific PCR technique with two different primer sets (Fig. 4e). Microsatellite analysis in the 3'-flanking region of porcine *C3* revealed three alleles of 211, 207 and 204 bp (amplicon C3-V), and these alleles corresponded to (T)<sub>21</sub>-, (T)<sub>17</sub>-, and (T)<sub>14</sub>-repeat units, respectively (Fig. 4f, allele 204 not shown).

Five of these polymorphic sites (at positions -44, 204 of intron 13, 1509, 3882 and +71) were segregating in the DUMI F2-resource population, and these alleles were coming from some Berlin Miniature Pig of the grandparent generation. All six polymorphic sites were segregating only in Thai native pigs, but no polymorphism was found among the 86 Duroc, German Landrace, Large White and Pietrain pigs. Allelic frequencies of the polymorphic sites in porcine *C3* gene among pig breeds are given in Table IV.

### 3.3. Association analysis

Means and standard deviation of the haemolytic complement activity of alternative and classical pathways were estimated (Tab. V). Only 3 SNPs





**Figure 4.** Polymorphism in the porcine *C3* gene including the 5'- and 3'-flanking regions (a); SSCP analysis at positions -384 and -44 bp in the 5'-flanking region (b); *TaqI*-PCR-RFLP test at position 204 in intron 13 (c); *Hsp92* I-PCR-RFLP test at position 1905 (d); allele specific PCR test at position 3882 (e); and SSCP analysis at position +71 bp in the 3'-flanking region (f).

and one microsatellite (nt 204 of intron 13, 1905, 3882, +71) were genotyped. These four alleles were segregating with two haplotype patterns that can be described [2] as follows: allele *C3*\*1, [IVS13+204T; 1905C; 3882G; c.5038+71(T)<sub>21</sub>] and allele *C3*\*2, [IVS13+204C; 1905A; 3882A; c.5038+71(T)<sub>17</sub>]. The numbers of pigs for each genotype class were 290, 109 and 17 for *C3*\*1/*C3*\*1, *C3*\*1/*C3*\*2 and *C3*\*2/*C3*\*2 genotypes respectively. Subsequently, the genotype *C3*\*2/*C3*\*2 was excluded from the analysis because of its low frequency and unequal distribution over parities and litters.

**Table IV.** Allele frequencies at the porcine *C3* gene in six genetic groups of pigs.

Position	Allele	Frequency					
		F2-DUMI	Duroc	German Landrace	Large White	Pietrain	Thai native pig
-384	G	1.00	1.00	1.00	1.00	1.00	0.99
-44	G	0.99	1.00	1.00	1.00	1.00	0.97
204 (intron 13)	T	0.75	1.00	1.00	1.00	1.00	0.82
1905	C	0.74	1.00	1.00	1.00	1.00	0.70
3882	G	0.74	1.00	1.00	1.00	1.00	0.82
+71(T) <sub>21</sub>	T <sub>21</sub>	0.76	1.00	1.00	1.00	0.00	0.71
+71(T) <sub>17</sub>	T <sub>17</sub>	0.24	0.00	0.00	0.00	0.00	0.12
+71(T) <sub>14</sub>	T <sub>14</sub>	0.00	0.00	0.00	0.00	0.00	0.17


**Table V.** Haemolytic complement activities (mean  $\pm$  SD) of alternative and classical pathways in DUMI resource population.

Blood sampling	Haemolytic complement activity (Unit $\cdot$ mL <sup>-1</sup> )	
	ACH50	CCH50
Time-1	53.64 $\pm$ 28.31	40.70 $\pm$ 23.13
Time-2	53.22 $\pm$ 28.43	48.23 $\pm$ 28.81
Time-3	55.05 $\pm$ 24.86	47.97 $\pm$ 29.19
Time-4	56.86 $\pm$ 30.13	59.08 $\pm$ 36.46
Time-5	62.31 $\pm$ 32.08	64.91 $\pm$ 35.71
Time-6	68.46 $\pm$ 47.63	59.79 $\pm$ 39.10
Time-7	69.75 $\pm$ 41.70	62.53 $\pm$ 35.58
Time-8	69.71 $\pm$ 37.42	60.99 $\pm$ 40.89

**Table VI.** Significance in analysis of variance of alternative (ACH50) and classical (CCH50) complement activity traits.

Traits	Test of fixed effects ( <i>P</i> value)			
	<i>C3</i> marker	Time	<i>C3</i> $\times$ time	Sire
ACH50	0.0255	< 0.0001	< 0.0001	< 0.0001
CCH50	0.0286	< 0.0001	0.0014	0.0151

Analysis of variance revealed that hemolytic complement activity of both alternative and classical pathways was significantly affected ( $P < 0.05$ ) by *C3* marker, time of blood sampling and their interaction, and by sire effect (Tab. VI). The profiles of the haemolytic complement activities between the different *C3*



**Figure 5.** Haemolytic complement activity profiles of alternative (a) and classical (b) pathways for pigs with *C3\*1/C3\*1* and *C3\*1/C3\*2* haplotypes.

genotypes were not parallel over time (Fig. 5). Complement activities in the alternative and classical pathway were  $5.8$  and  $6.9$  units  $\cdot$  mL<sup>-1</sup> higher for animals homozygous for the *C3\*1* allele than for *C3\*1/C3\*2* animals.

#### 4. DISCUSSION

Activated complement system plays an important role in killing and neutralization of microorganisms, and *C3* is a key molecule of this system. Therefore, *C3* can be regarded as a candidate gene for complement activities. The 5.1 kb sequence of full-length porcine *C3* gene has a 4983-bp ORF which coded for  $\alpha$ - and  $\beta$ -chain of *C3* protein. The porcine *C3* is a conserved gene and has high homology with the other mammalian species *C3* gene. Indeed, the putative promoter region contains one TATA-box and several cis-regulating

elements similar to that of the human *C3* promoter [8,21]. The variation of haemolytic complement activities before and after vaccination within the F2-DUMI resource population was found to be associated with *C3* markers. Although these *C3* markers were silent at the amino acids level, another nucleotide exchange in the 5'-flanking promoter region of the *C3* gene and in linkage disequilibrium with these *C3* variants might be responsible for its function. Animals with the more frequent *C3\*1/C3\*1* haplotype, present in Duroc and other commercial breeds, exhibited higher complement reactivity than those carrying the allele *C3\*2* coming from Berlin Miniature Pigs.

Genetic variation in *C3* affecting the complement activity has been observed in human [3], rabbit [11], and mouse [5]. These reports indicated that the genetic defect of *C3* gene caused low (or lack of) haemolytic complement activity and increased susceptibility to microorganisms. Phenotypic variation of haemolytic complement activity in pigs has been associated with effects due to vaccination, genomic variation at the swine leukocytes antigen locus (*SLA*) [16,20], breed [12], age as well as sire or/and dam [16,24].

Although the complement is a complex system and is controlled by many genes, the present work showed there was an association of *C3* and haemolytic complement activity – a defence mechanism active against many pathogenic agents – which reinforces the importance of *C3* as a candidate gene for natural resistance to microorganisms.

## ACKNOWLEDGEMENTS

We wish to express our great appreciation to Prof. Dr. S. Rattanarochart, Department of Animal Science, Faculty of Agriculture, Chiang Mai University, Thailand, for providing us with the blood samples. We would also like to thank the German research Council (DFG), and the German Academic Exchange Service (DAAD) for financial aid.

## REFERENCES

- [1] Ameratunga R., Winkelstein J.A., Brody L., Binns M., Cork L.C., Colombani P., Valle D., Molecular analysis of the third component of canine complement (*C3*) and identification of the mutation responsible for hereditary canine *C3* deficiency, *J. Immunol.* 160 (1998) 2824–2830.
- [2] Antonarakis S.E., Nomenclature Working Group, Recommendations for a nomenclature system for human gene mutations, *Hum. Mutat.* 11 (1998) 1–3.
- [3] Botto M., Fong K.Y., So A.K., Rudge A., Walport M.J., Molecular basis of hereditary *C3* deficiency, *J. Clin. Invest.* 86 (1990) 1158–1163.
- [4] Botto M., Fong K.Y., So A.K., Barlow R., Routier R., Morley B.J., Walport M.J., Homozygous hereditary *C3* deficiency due to a partial gene deletion, *Proc. Natl. Acad. Sci. USA* 89 (1992) 4957–4961.

- [5] Circolo A., Garnier G., Fukuda W., Wang X., Hidvegi T., Szalai A.J., Briles D.E., Volanakis J.E., Wetsel R.A., Colten H.R., Genetic disruption of the murine complement *C3* promoter region generates deficient mice with extrahepatic expression of *C3* mRNA, *Immunopharmacology* 42 (1999) 135–149.
- [6] de Bruijn M.H., Fey G.H., Human complement component *C3*: cDNA coding sequence and derived primary structure, *Proc. Natl. Acad. Sci. USA* 82 (1985) 708–712.
- [7] Fey G.H., Lundwall A., Wetsel R.A., Tack B.F., de Bruijn M.H., Domdey H., Nucleotide sequence of complementary DNA and derived amino acid sequence of murine complement protein *C3*, *Philos. Trans R. Soc. Lond. B. Biol. Sci.* 306 (1984) 333–344.
- [8] Fong K.Y., Botto M., Walport M.J., So A.K., Genomic organization of human complement component *C3*, *Genomics* 7 (1990) 579–586.
- [9] Hardge T., Köpke K., Reißmann M., Wimmers K., Maternal influences on litter size and growth in reciprocal crossed miniature pigs and Durocs, *Arch. Tierz. Dummerstorf* 42 (1999) 83–92.
- [10] Kawamura N., Singer L., Wetsel R.A., Colten H.R., Cis- and trans-acting elements required for constitutive and cytokine-regulated expression of the mouse complement *C3* gene, *Biochem. J.* 283 (1992) 705–712.
- [11] Komatsu M., Yamamoto K., Nakano Y., Nakazawa M., Ozawa A., Mikami H., Tomita M., Migita S., Hereditary *C3* hypocomplementemia in the rabbit, *Immunology* 64 (1988) 363–368.
- [12] Komatsu M., Yoshiwara S., Akita T., Immunological characteristics of the meishan pig: serum complement activity, serum *C3* level, immune response and skin structure, in: *Proceedings of the 6th World Congress on Genetics Applied to Livestock Production*, 11–16 January 1998, Vol. 26, University of New England, Armidale, pp. 323–326.
- [13] Kopf M., Abel B., Gallimore A., Carroll M., Bachmann M.F., Complement component *C3* promotes T-cell priming and lung migration to control acute influenza virus infection, *Nat. Med.* 8 (2002) 373–378.
- [14] Littell R.C., Henry P.R., Ammerman C.B., Statistical analysis of repeated measures data using SAS<sup>®</sup> procedures, *J. Anim. Sci.* 76 (1998) 1216–1231.
- [15] Liu C.C., Young J.D., A semiautomated microassay for complement activity, *J. Immunol. Methods* 114 (1988) 33–39.
- [16] Mallard B.A., Wilkie B.N., Kennedy B.W., Influence of major histocompatibility genes on serum hemolytic complement activity in miniature swine, *Am. J. Vet. Res.* 50 (1989) 359–363.
- [17] Misumi Y., Sohda M., Ikehara Y., Nucleotide and deduced amino acid sequence of rat complement *C3*, *Nucleic Acids Res.* 25 (1990) 2178.
- [18] Ren Y., Liao W.S., Transcription factor AP-2 functions as a repressor that contributes to the liver-specific expression of serum amyloid A1 gene, *J. Biol. Chem.* 276 (2001) 17770–17778.
- [19] Sahu A., Lambris J.D., Structure and biology of complement protein *C3*, a connecting link between innate and acquired immunity, *Immunol. Rev.* 180 (2001) 35–48.

- [20] Vaiman M., Hauptmann G., Mayer S., Influence of the major histocompatibility complex in the pig (SLA) on serum haemolytic complement levels, *J. Immunogenet.* 5 (1978) 59–65.
- [21] Vik D.P., Amiguet P., Moffat G.J., Fey M., Amiguet-Barras F., Wetsel R.A., Tack B.F., Structural features of the human *C3* gene: intron/exon organization, transcriptional start site, and promoter region sequence, *Biochemistry* 30 (1991) 1080–1085.
- [22] Wessels M.R., Butko P., Ma M., Warren H.B., Lage A.L., Carroll M.C., Studies of group B streptococcal infection in mice deficient in complement component *C3* or *C4* demonstrate an essential role for complement in both innate and acquired immunity, *Proc. Natl. Acad. Sci. USA* 92 (1995) 11490–11494.
- [23] Wilson D.R., Juan T.S., Wilde M.D., Fey G.H., Darlington G.J., A 58-base-pair region of the human *C3* gene confers synergistic inducibility by interleukin-1 and interleukin-6, *Mol. Cell Biol.* 10 (1990) 6181–6191.
- [24] Wimmers K., Lipperheide C., Ponsuksili S., Schmoll F., Hardge T., Petersen B., Schellander K., Haemolytic complement activity and C3c serum concentration in pigs, *Arch. Tierz. Dummerstorf* 42 (1999) 93–102.
- [25] Wimmers K., Mekchay S., Ponsuksili S., Hardge T., Schellander K., Polymorphic sites in exon 15 and 30 of the porcine *C3* gene, *Anim. Genet.* 32 (2001) 46–47.